



# DMC

*De Mathematics Competitions*

## Official Solutions

De Mathematics Competitions

3rd Annual

# DMC 10 B

Friday, October 28, 2022



This official solutions booklet gives at least one solution for each problem on this year's competition and shows that all problems can be solved without the use of a calculator. When more than one solution is provided, this is done to illustrate a significant contrast in methods. These solutions are by no means the only ones possible, nor are they necessarily superior to others the reader may devise.

People are encouraged to share copies of the problem booklet and official solutions with their friends and acquaintances to make the DMC Committee feel happy. All problems should be credited to the DMC (for example, "2020 DMC 10, Problem #21"). The publication, reproduction, or communication of the competition's problems or solutions for revenue-generating purposes is illegal.

Questions and complaints about this competition should be sent by private message to

**DeToasty3.**

The problems and solutions for this DMC 10 were prepared by the DMC 10 Editorial Board under the direction of

bronzetruck2016, cj13609517288, DankBasher619, dc495, DeToasty3, firebolt360, Hriship, john0512, NH14, nikenissan, pandabearcat, PhunsukhWangdu, pog, RedFlame2112, smartatmath, stayhomedomath, treemath, vsamc, YBSuburbanTea, & yusufsheikh2207

(Also, special thanks to GammaZero, kimi\_sun, Oxymoronic15, & richy for contributing problems, and to P\_Groudon & peace09 for helping with this solutions booklet!)

## Answer Key:

1. (D)	2. (C)	3. (E)	4. (C)	5. (E)
6. (E)	7. (D)	8. (B)	9. (D)	10. (A)
11. (E)	12. (D)	13. (A)	14. (C)	15. (E)
16. (C)	17. (E)	18. (C)	19. (D)	20. (C)
21. (D)	22. (D)	23. (B)	24. (E)	25. (D)

### Problem 1:

**(Oxymoronic15)** What is the value of  $\frac{2022! \cdot 2019!}{2020! \cdot 2021!}$ ?

- (A)  $\frac{1009}{1010}$     (B)  $\frac{2020}{2021}$     (C)  $\frac{2021}{2022}$     (D)  $\frac{1011}{1010}$     (E)  $\frac{2023}{2022}$

**Answer (D):**

The given expression is equal to  $\frac{2022!}{2021!} \cdot \frac{2019!}{2020!}$ . Note that  $\frac{2022!}{2021!} = 2022$  and  $\frac{2019!}{2020!} = \frac{1}{2020}$ , so our answer is  $2022 \cdot \frac{1}{2020} = \frac{2022}{2020} = \boxed{\text{(D)} \frac{1011}{1010}}$ . ■

### Problem 2:

**(DeToasty3)** When Amanda multiplies her favorite number by 3, subtracts the result from 14, and divides the result by 4, the resulting number will be Amanda's favorite number. What is Amanda's favorite number?

- (A) -14    (B) -7    (C) 2    (D) 7    (E) 14

**Answer (C):**

Let Amanda's favorite number be  $x$ . Then  $\frac{14 - 3x}{4} = x$ , so  $14 - 3x = 4x$ . Thus,  $14 = 7x$ , so  $x = \boxed{\text{(C)} 2}$ . ■

### Problem 3:

**(kimi\_sun)** Jeb has  $n$  toys. If he removes 3 toys,  $x\%$  of the toys remain. If, instead, he removes 6 toys,  $y\%$  of the toys remain. If  $x - y = 10$ , what is  $n$ ?

- (A) 6    (B) 10    (C) 24    (D) 27    (E) 30

**Answer (E):**

We get that  $(n - 3) - (n - 6) = 3$  is equal to 10% of  $n$ , so our answer is  $3 \cdot 10 = \boxed{\text{(E) } 30}$ . ■

**Problem 4:**

**(pog)** Ken writes 10 positive integers onto a sheet of paper. Joe then asks the following questions:

- How many of the numbers on your paper are less than 2?
- How many of the numbers on your paper are greater than 2?
- How many of the numbers on your paper are equal to 4?

Ken truthfully answers 3 to every question. What is the sum of the numbers on Ken's paper?


- (A) 21    (B) 22    (C) 23    (D) 24    (E) 25

**Answer (C):**

The only positive integer less than 2 is 1, so Ken wrote the numbers (1, 1, 1). The three numbers that Ken wrote that are greater than 2 are (4, 4, 4). Since Ken wrote 10 numbers, we get that  $10 - 3 - 3 = 4$  of Ken's numbers are equal to 2. Thus, Ken wrote (1, 1, 1, 2, 2, 2, 2, 4, 4, 4), so the requested sum is  $3 + 8 + 12 = \boxed{\text{(C) } 23}$ . ■

**Problem 5:**

**(pog)** Two congruent rectangles each with height 2 are stacked on top of each other. The top sheet is then rotated  $90^\circ$  about its center, resulting in the 12-sided polygon as shown.



If the area of this polygon is 30, what is the area of one of the rectangles?

- (A) 12    (B)  $6\sqrt{6}$     (C) 15    (D)  $5\sqrt{10}$     (E) 17

**Answer (E):**

Let the area of one rectangle be  $a$ . Then  $2a - 4 = 30$ , so  $a = \boxed{\text{(E) } 17}$ . ■

**Problem 6:**

**(GammaZero)** Using only the digits 2, 3 and 9, how many six-digit positive integers can be formed that are divisible by 6?

- (A) 27    (B) 35    (C) 36    (D) 80    (E) 81

**Answer (E):**

If a number is divisible by 6, it must be divisible by both 2 and 3. For a number to be even, its units digit must also be even, so the units digit of the number is 2.

Then, we want the sum of the digits of the number to be divisible by 3. Since 3 and 9 are both multiples of 3, the only thing that affects the sum of the digits (mod 3) is the number of twos in the number. We get that there are either 3 twos in the number or 6 twos in the number. For the first case, there are  $\binom{6}{3} \cdot 2^3 = 80$  numbers that can be formed, and for the second case, there is only 1 number that can be formed, namely 222222.

Thus, the number of six-digit positive integers that can be formed that are divisible by 6 is  $80 + 1 = \boxed{\text{(E) } 81}$ . ■

**Problem 7:**

**(DeToasty3)** Let  $a$ ,  $b$ , and  $c$  be consecutive positive integers, and let  $p$ ,  $q$ , and  $r$  also be consecutive positive integers, both not necessarily in order. Given that  $a \cdot p = 161$  and  $b \cdot q = 189$ , what is  $c \cdot r$ ?

- (A) 128    (B) 144    (C) 160    (D) 176    (E) 192

**Answer (D):**

The prime factorization of 161 is  $23 \cdot 7$ . Hence, since  $a$  and  $p$  are positive integers,  $\{a, p\}$  is either  $\{23, 7\}$  or  $\{161, 1\}$ . Since  $a$ ,  $b$ , and  $c$  are consecutive positive integers, any two of them differ in absolute value by 2 or less.

If  $\{a, p\} = \{161, 1\}$  and  $a = 161$  (without loss of generality), then  $159 \leq$

$b \leq 163$ . However, since  $b$  is an integer and  $b \cdot q = 189$ ,  $b$  is a divisor of 189, but no values of  $b$  in range are divisors of 189. Hence,  $\{a, p\} = \{161, 1\}$  is not possible.

Thus,  $\{a, p\} = \{23, 7\}$ . Without loss of generality, assume that  $a = 23$  and  $p = 7$ . Then,  $21 \leq b \leq 25$ . In this range, only  $b = 21$  is a divisor of 189. This implies that  $q = 9$ . Since  $\{a, b, c\}$  and  $\{p, q, r\}$  are both sets of three consecutive integers, the only possibility is  $c = 22$  and  $r = 8$ , implying that  $c \cdot r = \boxed{\text{(D) } 176}$ . ■

## Problem 8:

**(DeToasty3)** How many ways can the six variables in the equation

$$a + b + c = d + e + f + 2$$

be set equal to one of the numbers 1, 2, and 3 such that each of the three numbers is used by exactly two variables?

- (A) 9      (B) 18      (C) 24      (D) 27      (E) 36

**Answer (B):**

Since each value is used for exactly two variables,  $a + b + c + d + e + f = 2(1 + 2 + 3) = 12$ . With the given equation, it follows that  $d + e + f = 5$  and  $a + b + c = 7$ .

Since each variable is one of 1, 2, or 3 (with each being used twice), there are only a few possibilities:

- $\{a, b, c\} = \{3, 2, 2\}$  and  $\{d, e, f\} = \{3, 1, 1\}$
- $\{a, b, c\} = \{3, 3, 1\}$  and  $\{d, e, f\} = \{2, 2, 1\}$

Taking permutations of the sets into account, the number of ways is  $3 \cdot 3 + 3 \cdot 3 =$

$\boxed{\text{(B) } 18}$ . ■

## Problem 9:

**(DeToasty3)** Daniel is walking on a field, starting at his house. After walking 3 miles due north and 4 miles due west, Daniel arrives at the market. From the market, if Daniel walks  $n$  miles due east, Daniel will arrive at his school, which is the same distance from his house as it is from the market. What is  $n$ ?

- (A)  $\frac{9}{4}$       (B)  $\frac{21}{8}$       (C) 3      (D)  $\frac{25}{8}$       (E)  $\frac{25}{6}$

**Answer (D):**

Let the field be in the coordinate plane, where Daniel's house is the origin. Then, the market is at  $(-4, 3)$  and his school is at  $(n - 4, 3)$ . Since his school is equidistant from his house and from the market, by the distance formula

$$n = \sqrt{(n - 4)^2 + 3^2} \implies 8n = 25 \implies \boxed{\text{(D)} \frac{25}{8}}.$$

■

**Problem 10:**

**(pog)** Suppose that  $f(x) = 2x^2 + ax + b$  and  $g(x) = x^2 + cx + d$  intersect at  $x = 3$  and  $x = 4$ . If  $f(5) = 40$ , what is  $g(5)$ ?

- (A) 38    (B) 39    (C) 40    (D) 41    (E) 42

**Answer (A):**


We get that  $f(x) = g(x)$  at  $x = 3$  and  $x = 4$ , so

$$f(x) - g(x) = x^2 + (a - c)x + (b - d)$$

has roots 3 and 4. Hence,  $f(x) - g(x) = x^2 - 7x + 12$ . Thus,  $f(5) - g(5) = 5^2 - 7 \cdot 5 + 12$ . Substituting, we get  $40 - g(5) = 2$ , so  $g(5) = \boxed{\text{(A)} 38}$ . ■

**Problem 11:**

**(pog)** How many parallelograms are in the diagram below?



- (A) 18    (B) 31    (C) 36    (D) 39    (E) 40

**Answer (E):**

There are two cases to consider: if we do not include the topmost parallelogram, we are simply choosing a parallelogram from a grid of  $3 \times 3$  parallelograms. There are  $\binom{4}{2}$  ways to choose the vertical edges of the paral-

lelogram, and  $\binom{4}{2}$  ways to choose the horizontal edges of the parallelogram, for a total of  $\binom{4}{2} \cdot \binom{4}{2} = 36$  parallelograms that do not include the topmost parallelogram. Then, there are an additional 4 parallelograms that include the topmost parallelogram, for an answer of  $36 + 4 = \boxed{\text{(E) } 40}$ . ■

## Problem 12:

**(pog)** In a survey, each respondent is either a truth-teller or a liar and has a favorite number of either 2 or 5, but not both. Truth-tellers always tell the truth, and liars always lie. Each respondent answered two questions:

- Is your favorite number 2?
- Is your favorite number either 2 or 5?

If 38 liars have a favorite number of 2, 87 respondents answered “Yes” to the first question, and 106 respondents answered “No” to the second question, how many truth-tellers have a favorite number of 2?

- (A) 16    (B) 17    (C) 18    (D) 19    (E) 20

### Answer (D):

The second question must be true. If 106 respondents answered “No” to the second question, then there are 106 liars. Hence,  $106 - 38 = 68$  liars have a favorite number of 5. Consequently, 68 liars answered “Yes” to the first question, and the other  $87 - 68 = \boxed{\text{(D) } 19}$  people are the truth-tellers that have a favorite number of 2. ■

## Problem 13:

**(pog)** The number of positive divisors of the number  $(12!)^{26}$  can be written as

$$\underline{1} \underline{3} \underline{2} \underline{A} \underline{0} \underline{3} \underline{7} \underline{6} \underline{6} \underline{B},$$

where  $A$  and  $B$  are digits. What is the ordered pair  $(A, B)$ ?

- (A) (1, 7)    (B) (2, 6)    (C) (6, 3)    (D) (7, 6)    (E) (9, 8)

### Answer (A):

Note that  $(12!)^{26}$  is a perfect square (which always has an odd number of divisors), so  $B$  must be odd. This eliminates answer choices (B), (D), and (E).

As well,  $12!$  has exactly one factor of 11, so  $(12!)^{26}$  will have  $11^{26}$  in its

prime factorization. Thus, the number of positive divisors of  $(12!)^{26}$  will be divisible by  $26 + 1 = 27$ . Hence,  $1 + 3 + 2 + A + 0 + 3 + 7 + 6 + 6 + B \equiv 0 \pmod{9}$ , so  $A + B \equiv 8 \pmod{9}$ , eliminating answer choice (C). The only answer choice that works is **(A) (1, 7)**. ■

## Problem 14:

**(pog)** In a game, Bill has the numbers  $-1$ ,  $5$ , and  $10$ , while Ben has the numbers  $0$ ,  $2$ ,  $4$ ,  $6$ , and  $8$ . Each round, Bill and Ben randomly pick one of their own numbers, and the player with the higher number earns a point. The winner is the player with the most points after 3 rounds. Given that each player picks a different number each round, what is the probability that Bill wins?

- (A)  $\frac{1}{2}$     (B)  $\frac{11}{20}$     (C)  $\frac{3}{5}$     (D)  $\frac{2}{3}$     (E)  $\frac{3}{4}$

### Answer (C):

Note that Bill is guaranteed to win the round that he plays  $10$  and lose the round that he plays  $-1$ . Thus, in order to win the game, he must win the round where he plays  $5$  to get two points. This means that Ben must play  $0$ ,  $2$ , or  $4$  in the round where Bill plays  $5$ . By symmetry, the probability that Ben plays  $0$ ,  $2$ , or  $4$  is **(C)  $\frac{3}{5}$** . ■

## Problem 15:

**(pog)** Ayaka and Judo are each given a whole number greater than  $23$  and less than  $35$ . Ayaka is also told the units digit of Judo's number, while Judo forgets the units digit of his own number. They are told that both of their numbers are greater than  $23$  and less than  $35$ . Both of them know where each digit they remember is located in their two numbers, that Ayaka knows the units digit of Judo's number, and that they could have the same number.

- **Ayaka:** I don't know whether my number is less than your number.
- **Judo:** Oh, then I know whose number is larger.
- **Ayaka:** The positive difference between our numbers is  $5$ .

What is the sum of Ayaka and Judo's numbers?

- (A) 53    (B) 55    (C) 59    (D) 61    (E) 63



**Answer (E):**

We will go statement by statement.

**Statement 1:**

If Ayaka is unsure, then the units digit of Judo's number must be 4. If the units digit of Judo's number was not 4, then Ayaka could just determine Judo's number and figure out if her number is greater than Judo's number or not. (If the two numbers are equal, the answer would be no.)

As well, Ayaka's number cannot be 34, because then she would know her number must not be less (i.e. must be greater than or equal to) than Judo's number regardless of if it was 24 or 34, which contradicts her statement.

**Statement 2:**

Judo now knows that his units digit is 4 and that Ayaka's number is not equal to 34.

**Case 1:** Judo remembers the tens digit 2.

If Judo remembered the tens digit 2, then he now knows his number is 24. However, this doesn't corroborate with his statement, as he knows Ayaka's number could still be 24, so he could not determine for certain whose number is larger.

**Case 2:** Judo remembers the tens digit 3.

If Judo remembered the tens digit 3, then he now knows his number is 34. He would then know that his number is larger than Ayaka's as he knows her number is not 34.

Hence, Judo's number must be 34.

**Statement 3:**

Based on this statement, we find that Ayaka's number is 29, so our answer is  $29 + 34 = \boxed{\text{(E) } 63}$ . ■

**Problem 16:**

**(pog)** The number  $2022 \cdot 10^{21}$  is written on a piece of paper. One day, Katherine repeatedly erases the number on the paper, divides it by 512, and writes the result.

This process continues until the number on the paper is less than 1. How many times did Katherine divide the number on the paper by 512?

- (A) 7      (B) 8      (C) 9      (D) 10      (E) 11

**Answer (C):**

Recall that  $2^9 = 512$ . Suppose Katherine divides the number on the paper  $k$  times. Then, the number she gets after the  $(k - 1)$ th division must be greater than 1, and then the next number is less than 1. Hence,  $\frac{2022 \cdot 10^{21}}{2^{9(k-1)}} > 1$  and  $\frac{2022 \cdot 10^{21}}{2^{9k}} < 1$ . Combining these two inequalities implies

$$2^{9(k-1)} < 2022 \cdot 10^{21} < 2^{9k}.$$

To find a lower bound on the expression, note that  $2022 > 1024 = 2^{10}$  and  $10 > 8 = 2^3$ . Hence,

$$2022 \cdot 10^{21} > 2^{10} \cdot (2^3)^{21} = 2^{73} > 2^{72}.$$

To find an upper bound on the expression, note that  $2022 < 2048 = 2^{11}$  and  $10^3 = 1000 < 1024 = 2^{10}$ . Hence,

$$2022 \cdot 10^{21} < 2^{11} \cdot (10^3)^7 < 2^{11} \cdot (2^{10})^7 = 2^{81}.$$

Combining the two inequalities,

$$2^{72} < 2022 \cdot 10^{21} < 2^{81}.$$

Thus, the value of  $k$  is **(C) 9**. ■

## Problem 17:

**(DeToasty3)** A line passes through the point  $A(5, 4)$  and has slope  $-8$ . A second line passes through the point  $B(1, 6)$  and intersects the first line at a point  $C$ , equidistant from  $A$  and  $B$ . What is the slope of the second line?

- (A)  $\frac{1}{3}$       (B)  $\frac{2}{5}$       (C)  $\frac{1}{2}$       (D)  $\frac{6}{11}$       (E)  $\frac{4}{7}$

**Answer (E):**

Let  $\ell_1$  and  $\ell_2$  be the first two lines.

Since  $C$  is equidistant from  $A$  and  $B$ , it lies on the perpendicular bisector

of  $\overline{AB}$ , which will be defined as line  $\ell_3$ . Line  $\ell_1$  is given by the equation  $y - 4 = -8(x - 5)$ . If  $M$  is the midpoint of  $\overline{AB}$ , then  $M = (3, 5)$ . In addition,  $\ell_3$  passes through  $M$  and is perpendicular to  $AB$ . Since line  $AB$  has slope  $-\frac{1}{2}$ , so the equation for  $\ell_3$  is  $y - 5 = 2(x - 3)$ .

By definition,  $\ell_1$  and  $\ell_3$  intersect at  $C$ , where the solution to the system is  $(x, y) = (\frac{9}{2}, 8)$ . Then, since  $\ell_2$  contains  $B$  and  $C$ , its slope is

$$\frac{8 - 6}{\frac{9}{2} - 1} = \boxed{\text{(E)} \frac{4}{7}}.$$

## Problem 18:


**(DeToasty3)** A rectangle has perimeter 36. The rectangle is split into three smaller rectangles with dimensions 9-by-4, 6-by-5, and  $m$ -by- $n$ . What is  $m + n$ ?

- (A) 6      (B) 7      (C) 8      (D) 9      (E) 10


### Answer (C):

Consider how three smaller rectangles can be placed to form a larger rectangle. Suppose two rectangles touch and without loss of generality, suppose they touch horizontally.

If there are no horizontal sides that line up as shown in the diagrams below, it is impossible to add another rectangle to the figure to form a bigger rectangle (since there are two spots where the dimensions need to be fixed to form the rectangle).




It then follows that the formation of the rectangles will appear like one of the two diagrams below:



The second formation is impossible with the given dimensions of the smaller rectangles, since it would require each rectangle to share a common dimension.

In the first formation, it can be seen that two of the rectangles (label these  $A$  and  $B$ ) share a common dimension. One of these two rectangles must be  $m \times n$ , as the  $9 \times 4$  and  $6 \times 5$  do not share any dimensions. Without loss of generality, let  $A$  be the  $m \times n$  rectangle, and let  $m$  be the shared dimension. See point labels:



Rectangle  $CDGI$  is either the  $9 \times 4$  one or the  $6 \times 5$ . Since the perimeter of  $CDEF$  is 36, it follows that  $CD + CF = 18$ .

If  $CDGI$  is the  $6 \times 5$  rectangle, then  $CD + CI = 11$ , implying  $m = 7$ . However, the  $9 \times 4$  rectangle is forced to be  $EGJH$ , which does not have a dimension of 7. Thus,  $CDGI$  cannot be the  $6 \times 5$  rectangle.

Hence,  $CDGI$  is the  $9 \times 4$  rectangle, implying  $m = 5$ . Thus,  $EGJH$  is the  $6 \times 5$  rectangle, implying that  $HE = JG = 6$ . Since  $IG > JG$ , it follows that  $IG = 9$  and  $CI = 4$  (and not the other way around). Thus,  $n = 9 - 6 = 3$ , implying  $m + n = \boxed{\text{(C) } 8}$ . ■

## Problem 19:

**(DeToasty3)** There are 2022 members in a math tournament, where 999 members are girls, and the rest are boys. The members are split into 674 groups of 3. For every two members in a group, if at least one of them is a girl, they shake hands. Otherwise, they do not. A member is *handy* if they shake hands with both members in their group. Let  $N$  be the maximum number of handy members in the math tournament. What is the sum of the digits of  $N$ ?

- (A) 6      (B) 15      (C) 16      (D) 22      (E) 23

**Answer (D):**

Consider each outcome of a group. If three people are girls, then all three of them are social. If two people are girls and one person is a boy, then all three of them are social. If one person is a girl and two people are boys, then one person is social. If three people are boys, then none of them are social. We want to maximize the number of groups where all three are social by using as few girls as possible, or two girls in a group. Thus, we have 499 groups with two girls and one boy, one group with one girl and two boys, and the other groups with three boys. giving  $499 \cdot 3 + 1 = 1498$ , so our answer is  $1 + 4 + 9 + 8 = \boxed{\text{(D) } 22}$ . ■

**Problem 20:**

**(richy)** Four people are in a tournament where every person duels each other person exactly once. Every duel ends in one person winning and the other losing (i.e., no ties). After the tournament, each person counts the number of wins they have and adds one, and then their numbers are multiplied. What is the smallest possible resulting product?


- (A) 16    (B) 18    (C) 24    (D) 27    (E) 32

**Answer (C):**

Let the wins for people 1, 2, 3, 4 be  $(w, x, y, z)$ , where  $w \leq x \leq y \leq z$ . We must have that  $w + x + y + z = 6$ . Note that two people cannot win 3 games, so we can limit the cases. Eventually, we get that only the ordered tuples  $(0, 1, 2, 3)$  and  $(0, 2, 2, 2)$  work. Clearly the first ordered tuple gives a smaller resulting product than the second ordered tuple does, so the answer is  $(0 + 1)(1 + 1)(2 + 1)(3 + 1) = \boxed{\text{(C) } 24}$ . ■

**Problem 21:**


**(DeToasty3)** In the figure below, six congruent rectangles are glued to each of the sides of a regular hexagon with side length 2, and six of the vertices of the rectangles are connected to form a regular hexagon with side length 4. The length of a side of one of the rectangles not equal to 2 can be written as  $\sqrt{m} - \sqrt{n}$ , where  $m$  and  $n$  are positive integers. What is  $m + n$ ?



- (A) 15    (B) 16    (C) 17    (D) 18    (E) 19

**Answer (D):**

See the diagram below for point labels, where  $O$  is the center of both hexagons and  $D$  is the foot of the altitude from  $O$  to line  $AB$ .



Let  $s$  be the side length of each of the rectangles other than 2. Then,  $AB = s$ . By a property of regular hexagons, the distance from its center to any of its vertices is the same as the hexagon's side length. Thus,  $OA = 2$  and  $OB = 4$ . Clearly,  $\angle OAB = \angle OAC + \angle CAB = 60^\circ + 90^\circ = 150^\circ$ . Hence,  $\angle OAD = 30^\circ$ . It follows that  $\triangle AOD$  is a 30-60-90 triangle. It immediately follows that  $OD = 1$  and  $DA = \sqrt{3}$ . Then, by Pythagorean Theorem on  $\triangle ODB$ ,

$$1^2 + (s + \sqrt{3})^2 + 4^2 \implies s + \sqrt{3} = \pm\sqrt{15} \implies s = \sqrt{15} - \sqrt{3}.$$

Thus,  $m + n = \boxed{\text{(D) } 18}$ . ■

## Problem 22:

**(HrishiP)** Let  $P(x)$  be a polynomial with degree 3 and roots  $r$ ,  $s$ , and  $t$  with sum 25 such that the coefficient of the  $x^3$  term is 1, and

$$(r + s)(s + t)(t + r) = 2500 \quad \text{and} \quad \left(\frac{1}{r} + \frac{1}{s}\right)\left(\frac{1}{s} + \frac{1}{t}\right)\left(\frac{1}{t} + \frac{1}{r}\right) = 100$$

are satisfied. If the constant term of  $P(x)$  is positive, the value of  $P(1)$  is equal to  $\frac{m}{n}$  for relatively prime positive integers  $m$  and  $n$ . What is  $m + n$ ?

(A) 406    (B) 407    (C) 408    (D) 409    (E) 410

**Answer (D):**

Let  $P(x) = x^3 - 25x^2 + bx + c$ . The desire is to find the value of  $1 - 25 + b + c$ . By Vieta's,  $r + s + t = -\frac{-25}{1} = 25$ . Then,

$$(25 - t)(25 - r)(25 - s) = 420.$$

We note that this is telling us  $P(25) = 420$ , or

$$\begin{aligned} 25^3 - 25(25^2) + 25b + c &= 2500 \\ 25b + c &= 2500. \end{aligned}$$

Next, we see that the second equation is equivalent to

$$\begin{aligned} \left(\frac{r+s}{rs}\right)\left(\frac{s+t}{st}\right)\left(\frac{t+r}{tr}\right) &= \frac{(r+s)(s+t)(t+r)}{(rst)^2} \\ &= \frac{2500}{(rst)^2} \\ &= \frac{2500}{c^2}, \end{aligned}$$

where the last equality is from Vieta's. Then,  $\frac{2500}{c^2} = 100$  or  $c^2 = 25$ . If we solve the system

$$25b + c = 2500, \quad c^2 = 25,$$

we get solutions  $(b, c) = \left(\frac{501}{5}, -5\right), \left(\frac{499}{5}, 5\right)$ . However,  $c > 0$ , so the last solution is the only one that works. Then,

$$P(1) = 1 - 25 + \frac{499}{5} + 5 = \frac{404}{5},$$

so  $m + n = \boxed{\text{(D) 409}}$ . ■


**Problem 23:**

**(DeToasty3)** Let parallelogram  $ABCD$  have  $BC = 5$ ,  $\angle ABC < 90^\circ$ , and  $\angle ACB > 90^\circ$ . Let line  $AD$  and side  $\overline{CD}$  intersect the circle passing through  $A$ ,  $B$ , and  $C$  at  $P \neq A$  and  $Q \neq C$ , respectively. If  $CP = 10$  and  $CQ = 4$ , what is  $AP$ ?

(A)  $\frac{48}{7}$     (B) 7    (C)  $\frac{36}{5}$     (D)  $\frac{15}{2}$     (E) 8

**Answer (B):**

Let  $\omega$  be the circumcircle of  $\triangle ABC$ . Since it is given that  $\angle ACB$  is obtuse, the center of  $\omega$  would be outside  $\triangle ABC$ . By constructing parallelogram  $ABCD$  from the locations of  $A$ ,  $B$ , and  $C$ , it is clear that  $D$  will be outside  $\omega$  and  $Q$  will be between  $C$  and  $D$ . Thus,  $A$  will be between  $P$  and  $D$ .



Since  $ACBP$  is a quadrilateral inscribed in a circle with  $\overline{AP} \parallel \overline{BC}$ ,  $ACBP$  is an isosceles trapezoid. Thus,  $AB = CP = 10$ . Then, by the parallelogram,  $DC = AB = 10$  and  $AD = BC = 5$ . With  $CQ = 4$ , it follows that  $QD = 6$ . Then, by Power of a Point at  $D$ ,

$$DA \cdot DP = DQ \cdot DC \implies 5 \cdot DP = 6 \cdot 10 \implies DP = 12.$$

Hence,  $AP = \boxed{\text{(B) } 7}$ . ■

**Problem 24:**

**(john0512)** For each positive integer  $n$ , let  $f_1(n) = n!$ , and for  $k \geq 2$ , let

$$f_k(n) = f_{k-1}(1) \cdot f_{k-1}(2) \cdot \dots \cdot f_{k-1}(n).$$

Let  $N$  be the largest integer such that  $f_4(10)$  is divisible by  $2^N$ . What is the sum of the digits of  $N$ ?

- (A) 4      (B) 15      (C) 16      (D) 20      (E) 21

**Answer (E):**

By taking  $n - 1$  instead of  $n$  in the function,

$$f_k(n - 1) = f_{k-1}(1) \cdot f_{k-1}(2) \cdot \dots \cdot f_{k-1}(n - 1),$$

so the given relation can be rewritten as

$$f_k(n) = f_k(n - 1) \cdot f_{k-1}(n). \quad (*)$$

Let  $\nu_2(x)$  denote the largest integer  $a$  such that  $2^a$  divides  $x$ . Then,  $\nu_2(f_4(10)) = N$  and by  $(*)$ ,



$$v_2(f_k(n)) = v_2(f_k(n-1)) + v_2(f_{k-1}(n)). \quad (\star)$$

By the definition of a factorial,  $f_1(n) = f_1(n-1) \cdot n$ , implying that  $v_2(f_1(n)) = v_2(f_1(n-1)) + v_2(n)$ . Using the base case of  $v_2(f_k(1)) = 0$  for all  $k \geq 1$  and  $(\star)$ , a recursion table can be filled out for  $v_2(f_k(n))$ :

$k \backslash n$	1	2	3	4	5	6	7	8	9	10
1	0	1	1	3	3	4	4	7	7	8
2	0	1	2	5	8	12	16	23	30	38
3	0	1	3	8	16	28	44	67	97	135
4	0	1	4	12	28	56	100	167	264	399

Thus,  $N = 399$ , so our answer is **(E) 21**. ■

## Problem 25:

**(stayhomedomath)** In acute  $\triangle ABC$  with  $AC < BC$ , the perpendicular bisector of  $\overline{AB}$  meets lines  $AB$ ,  $BC$ , and  $AC$  at  $D$ ,  $E$ , and  $F$ , respectively. If  $AD = 5$ ,  $BE = 13$ , and the area of  $\triangle ABC$  is 14 units greater than that of  $\triangle ADF$ , what is  $AF^2$ ?

- (A) 650      (B) 701      (C) 754      (D) 809      (E) 866

### Answer (D):

Henceforth, let the area of a polygon  $\mathcal{P}$  be denoted by  $[\mathcal{P}]$ . We shall first explain the setup of the configuration. Clearly,  $D$  is the midpoint of  $\overline{AB}$ .


Let  $\ell_1$ ,  $\ell_2$ , and  $\ell_3$  be the lines perpendicular to  $\overline{AB}$  passing through  $A$ ,  $D$ , and  $B$ , respectively. Then, in order for  $\triangle ABC$  to be acute,  $C$  must lie between  $\ell_1$  and  $\ell_3$ ; and since  $AC < BC$ ,  $C$  lies between  $\ell_1$  and  $\ell_2$ . As  $\ell_2$  is the perpendicular bisector of  $\overline{AB}$ , it follows that  $E$  lies in between  $D$  and  $F$ , with  $E$  on  $\overline{BC}$  and  $F$  on the extension of  $\overline{AC}$  past  $C$ .

Since  $AD = BD = 5$ ,  $DE = 12$  by the Pythagorean Theorem, and  $[BDE] = \frac{1}{2} \cdot 5 \cdot 12 = 30$ . Consequently,

$$[ABC] - [ADF] = ([BDE] + [ACED]) - ([CEF] + [ACED]) = 30 - [CEF],$$

and setting this equal to 14 gives  $[CEF] = 16$ .

Now, extend  $\overline{AE}$  past  $E$  to meet  $\overline{BF}$  at  $P$ . By symmetry,  $[ADE] = [BDE] = 30$  and  $[CEF] = [PEF] = 16$ ; accordingly denote  $[ACE] = [BPE] = K$ .



By area ratios,

$$\frac{[AEF]}{[PEF]} = \frac{AE}{PE} = \frac{[ABE]}{[PBE]} \implies \frac{K+16}{16} = \frac{60}{K}.$$

Cross-multiplying, we have that

$$K^2 + 16K - 960 = (K - 24)(K + 40) = 0 \implies K = 24.$$

Hence,  $[ADF] = 30 + 24 + 16 = 70$ , and  $DF = \frac{2[ADF]}{AD} = 28$ . Therefore,  $AF^2 = 5^2 + 28^2 = \boxed{\text{(D) } 809}$ . ■